

SPACE-TIME CONTINUOUS ANALYSIS OF WAVEFORM RELAXATION FOR THE HEAT EQUATION*

MARTIN J. GANDER[†] AND ANDREW M. STUART[‡]

Abstract. Waveform relaxation algorithms for partial differential equations (PDEs) are traditionally obtained by discretizing the PDE in space and then splitting the discrete operator using matrix splittings. For the semidiscrete heat equation one can show linear convergence on unbounded time intervals and superlinear convergence on bounded time intervals by this approach. However, the bounds depend in general on the mesh parameter and convergence rates deteriorate as one refines the mesh.

Motivated by the original development of waveform relaxation in circuit simulation, where the circuits are split in the physical domain into subcircuits, we split the PDE by using overlapping domain decomposition. We prove linear convergence of the algorithm in the continuous case on an infinite time interval, at a rate depending on the size of the overlap. This result remains valid after discretization in space and the convergence rates are robust with respect to mesh refinement. The algorithm is in the class of waveform relaxation algorithms based on overlapping multisplittings. Our analysis quantifies the empirical observation by Jeltsch and Pohl [*SIAM J. Sci. Comput.*, 16 (1995), pp. 40–49] that the convergence rate of a multisplitting algorithm depends on the overlap.

Numerical results are presented which support the convergence theory.

Key words. waveform relaxation, domain decomposition, overlapping Schwarz, multisplitting

AMS subject classifications. 65M55, 65M12, 65M15, 65Y05

PII. S1064827596305337

1. Introduction. The basic ideas of waveform relaxation were introduced in the late 19th century by Picard [18] and Lindelöf [11] to study initial value problems. There has been much recent interest in waveform relaxation as a practical parallel method for the solution of stiff ordinary differential equations (ODEs) after the publication of a paper by Lelarasmee, Ruehli, and Sangiovanni–Vincentelli [10] in the area of circuit simulation.

There are two classical convergence results for waveform relaxation algorithms for ODEs: (i) for linear systems of ODEs on unbounded time intervals one can show linear convergence of the algorithm under some dissipation assumptions on the splitting [15], [14], [4], [9]; (ii) for nonlinear systems of ODEs (including linear ones) on bounded time intervals one can show superlinear convergence assuming a Lipschitz condition on the splitting function [15], [1], [3]. For classical relaxation methods (Jacobi, Gauss–Seidel, SOR) the above convergence results depend on the discretization parameter if the ODE arises from a PDE which is discretized in space. The convergence rates deteriorate as one refines the mesh.

Jeltsch and Pohl propose in [9] a multisplitting algorithm with overlap, generalizing the elliptic analysis of O’Leary and White in [17] to the parabolic case. They prove results (i) and (ii) for their algorithm, but the convergence rates are mesh-dependent. However, they show numerically that increasing the overlap accelerates the convergence of the waveform relaxation algorithm. We quantify their numerical

*Received by the editors June 17, 1996; accepted for publication (in revised form) March 4, 1997; published electronically July 26, 1998.

<http://www.siam.org/journals/sisc/19-6/30533.html>

[†]Department of Computer Science, Stanford University, Room 286, Gates Building 2B, Stanford, CA 94305-2140 (mgander@sccm.stanford.edu).

[‡]SCCM Program, Division of Mechanics and Computation, Durand 257, Stanford University, Stanford, CA 94305-4040 (stuart@sccm.stanford.edu).

results by formulating the waveform relaxation algorithm at the space-time continuous level using overlapping domain decomposition; this approach was motivated by the work of Bjørhus [2]. We show linear convergence of this algorithm on unbounded time intervals at a rate depending on the size of the overlap. This is an extension of the first classical convergence result (i) for waveform relaxation from ODEs to PDEs. Discretizing the algorithm, the size of the physical overlap corresponds to the overlap of the multisplitting algorithm analyzed by Jeltsch and Pohl. We show furthermore that the convergence rate is robust with respect to mesh refinement, provided the physical overlap is held constant during the refinement process.

Giladi and Keller [8] study superlinear convergence of domain decomposition algorithms for the convection diffusion equation on bounded time intervals, hence generalizing the second classical waveform relaxation result (ii) from ODEs to PDEs.

It is interesting to note that, using multigrid to formulate a waveform relaxation algorithm, Lubich and Osterman prove in [13] linear convergence for the heat equation independent of the mesh parameter.

In section 2 we consider a decomposition of the domain into two subdomains. This section is mainly for illustrative purposes, since the analysis can be performed in great detail. In section 3 we generalize the results to an arbitrary number of subdomains. In section 4 we show numerical experiments which confirm the convergence results.

Although the analysis presented is restricted to the one-dimensional heat equation, the techniques applied in the proofs are more general. Future work successfully applies these techniques to higher-dimensional problems and to nonlinear parabolic equations.

2. Two subdomains.

2.1. Continuous case. Consider the one-dimensional heat equation on the interval $[0, L]$,

$$(2.1) \quad \begin{aligned} \frac{\partial u}{\partial t} &= \frac{\partial^2 u}{\partial x^2} + f(x, t), & 0 < x < L, \quad t > 0, \\ u(0, t) &= g_1(t), & t > 0, \\ u(L, t) &= g_2(t), & t > 0, \\ u(x, 0) &= u_0(x), & 0 \leq x \leq L, \end{aligned}$$

where we assume $f(x, t)$ to be bounded on the domain $[0, L] \times [0, \infty)$ and uniformly Hölder continuous on each compact subset of the domain. We assume furthermore that the initial data $u_0(x)$ and the boundary data $g_1(t), g_2(t)$ are piecewise continuous. Then (2.1) has a unique bounded solution [5]. We consider in the following functions in $L^\infty := L^\infty(\mathbb{R}^+; \mathbb{R})$ with the infinity norm

$$\|f(\cdot)\|_\infty := \sup_{t>0} |f(t)|.$$

The maximum principle, and a corollary thereof, establishing the steady state solution as a bound on the solution of the heat equation are instrumental in our analysis.

THEOREM 2.1 (maximum principle). *The solution $u(x, t)$ of the heat equation (2.1) with $f(x, t) \equiv 0$ attains its maximum and minimum either on the initial line $t = 0$ or on the boundary at $x = 0$ or $x = L$. If $u(x, t)$ attains its maximum in the interior, then $u(x, t)$ must be constant.*

Proof. The proof can be found in [21]. □

COROLLARY 2.2. *The solution $u(x, t)$ of the heat equation (2.1) with $f(x, t) \equiv 0$ and $u_0 \equiv 0$ satisfies the inequality*

$$(2.2) \quad \|u(x, \cdot)\|_\infty \leq \frac{L-x}{L} \|g_1(\cdot)\|_\infty + \frac{x}{L} \|g_2(\cdot)\|_\infty, \quad 0 \leq x \leq L.$$

Proof. Consider \tilde{u} solving

$$(2.3) \quad \begin{aligned} \frac{\partial \tilde{u}}{\partial t} &= \frac{\partial^2 \tilde{u}}{\partial x^2}, & 0 < x < L, \quad t > 0, \\ \tilde{u}(0, t) &= \|g_1(\cdot)\|_\infty, & t > 0, \\ \tilde{u}(L, t) &= \|g_2(\cdot)\|_\infty, & t > 0, \\ \tilde{u}(x, 0) &= \frac{L-x}{L} \|g_1(\cdot)\|_\infty + \frac{x}{L} \|g_2(\cdot)\|_\infty, & 0 \leq x \leq L. \end{aligned}$$

The solution \tilde{u} of (2.3) does not depend on t and is given by the steady state solution

$$\tilde{u}(x) = \frac{L-x}{L} \|g_1(\cdot)\|_\infty + \frac{x}{L} \|g_2(\cdot)\|_\infty.$$

By construction, we have $\tilde{u}(x) - u(x, t) \geq 0$ at $t = 0$, and on the boundary $x = 0$ and $x = L$. Since $\tilde{u} - u$ is in the kernel of the heat operator, we have by the maximum principle for the heat equation $\tilde{u}(x) - u(x, t) \geq 0$ on the whole domain $[0, L]$. Hence

$$u(x, t) \leq \frac{L-x}{L} \|g_1(\cdot)\|_\infty + \frac{x}{L} \|g_2(\cdot)\|_\infty.$$

Likewise $\tilde{u}(x) + u(x, t) \geq 0$ at $t = 0$, $x = 0$, and $x = L$, and is in the kernel of the heat operator. Hence

$$u(x, t) \geq - \left(\frac{L-x}{L} \|g_1(\cdot)\|_\infty + \frac{x}{L} \|g_2(\cdot)\|_\infty \right).$$

Therefore we have

$$|u(x, t)| \leq \frac{L-x}{L} \|g_1(\cdot)\|_\infty + \frac{x}{L} \|g_2(\cdot)\|_\infty.$$

Now the right-hand side does not depend on t , so we can take the supremum over t , which leads to the desired result. \square

To obtain a waveform relaxation algorithm, we decompose the domain $\Omega = [0, L] \times [0, \infty)$ into two overlapping subdomains $\Omega_1 = [0, \beta L] \times [0, \infty)$ and $\Omega_2 = [\alpha L, L] \times [0, \infty)$ where $0 < \alpha < \beta < 1$ as given in Figure 2.1. The solution $u(x, t)$ of (2.1) can now be obtained from the solutions $v(x, t)$ on Ω_1 and $w(x, t)$ on Ω_2 , which satisfy the equations

$$(2.4) \quad \begin{aligned} \frac{\partial v}{\partial t} &= \frac{\partial^2 v}{\partial x^2} + f(x, t), & 0 < x < \beta L, \quad t > 0, \\ v(0, t) &= g_1(t), & t > 0, \\ v(\beta L, t) &= w(\beta L, t), & t > 0, \\ v(x, 0) &= u_0(x), & 0 \leq x \leq \beta L, \end{aligned}$$

and

$$(2.5) \quad \begin{aligned} \frac{\partial w}{\partial t} &= \frac{\partial^2 w}{\partial x^2} + f(x, t), & \alpha L < x < L, \quad t > 0, \\ w(\alpha L, t) &= v(\alpha L, t), & t > 0, \\ w(L, t) &= g_2(t), & t > 0, \\ w(x, 0) &= u_0(x), & \alpha L \leq x \leq L. \end{aligned}$$




FIG. 2.1. Decomposition into two overlapping subdomains.

First note that $v = u$ on Ω_1 and $w = u$ on Ω_2 are solutions to (2.4) and (2.5). Uniqueness follows from our analysis of a Schwarz-type iteration introduced for elliptic problems in [19] and further studied in [12] and [6]. We get

$$\begin{aligned} \frac{\partial v^{k+1}}{\partial t} &= \frac{\partial^2 v^{k+1}}{\partial x^2} + f(x, t), & 0 < x < \beta L, \quad t > 0, \\ v^{k+1}(0, t) &= g_1(t), & t > 0, \\ v^{k+1}(\beta L, t) &= w^k(\beta L, t), & t > 0, \\ v^{k+1}(x, 0) &= u_0(x), & 0 \leq x \leq \beta L, \end{aligned}$$

and

$$\begin{aligned} \frac{\partial w^{k+1}}{\partial t} &= \frac{\partial^2 w^{k+1}}{\partial x^2} + f(x, t), & \alpha L < x < L, \quad t > 0, \\ w^{k+1}(\alpha L, t) &= v^k(\alpha L, t), & t > 0, \\ w^{k+1}(L, t) &= g_2(t), & t > 0, \\ w^{k+1}(x, 0) &= u_0(x), & \alpha L \leq x \leq L. \end{aligned}$$

Let $d^k(x, t) := v^k(x, t) - v(x, t)$ and $e^k(x, t) := w^k(x, t) - w(x, t)$ and consider the error equations

$$(2.6) \quad \begin{aligned} \frac{\partial d^{k+1}}{\partial t} &= \frac{\partial^2 d^{k+1}}{\partial x^2}, & 0 < x < \beta L, \quad t > 0, \\ d^{k+1}(0, t) &= 0, & t > 0, \\ d^{k+1}(\beta L, t) &= e^k(\beta L, t), & t > 0, \\ d^{k+1}(x, 0) &= 0, & 0 \leq x \leq \beta L, \end{aligned}$$

and

$$(2.7) \quad \begin{aligned} \frac{\partial e^{k+1}}{\partial t} &= \frac{\partial^2 e^{k+1}}{\partial x^2}, & \alpha L < x < L, \quad t > 0, \\ e^{k+1}(\alpha L, t) &= d^k(\alpha L, t), & t > 0, \\ e^{k+1}(L, t) &= 0, & t > 0, \\ e^{k+1}(x, 0) &= 0, & \alpha L \leq x \leq L. \end{aligned}$$

The following lemma establishes convergence of the Schwarz iteration on the interfaces of the subdomains in L^∞ . Using the maximum principle convergence in the interior follows.

LEMMA 2.3. *On the interfaces $x = \alpha L$ and $x = \beta L$ the error of the Schwarz iteration decays at the rate*

$$(2.8) \quad \|d^{k+2}(\alpha L, \cdot)\|_\infty \leq \frac{\alpha(1-\beta)}{\beta(1-\alpha)} \|d^k(\alpha L, \cdot)\|_\infty,$$

$$(2.9) \quad \|e^{k+2}(\beta L, \cdot)\|_\infty \leq \frac{\alpha(1-\beta)}{\beta(1-\alpha)} \|e^k(\beta L, \cdot)\|_\infty.$$

Proof. By Corollary 2.2 we have

$$(2.10) \quad \|d^{k+2}(x, \cdot)\|_\infty \leq \frac{x}{\beta L} \|e^{k+1}(\beta L, \cdot)\|_\infty \quad \forall x \in [0, \beta L]$$

and

$$(2.11) \quad \|e^{k+1}(x, \cdot)\|_\infty \leq \frac{L-x}{(1-\alpha)L} \|d^k(\alpha L, \cdot)\|_\infty \quad \forall x \in [\alpha L, L].$$

Evaluating (2.11) at $x = \beta L$ and (2.10) at $x = \alpha L$ and combining the two, we obtain inequality (2.8). Inequality (2.9) is obtained similarly. \square

For any function $g(\cdot, t)$ in $L^\infty([a, b], L^\infty)$ we introduce the norm

$$\|g(\cdot, \cdot)\|_{\infty, \infty} := \sup_{a \leq x \leq b} \|g(x, \cdot)\|_\infty.$$

THEOREM 2.4. *The Schwarz iteration for the heat equation with two subdomains converges in $L^\infty([a, b], L^\infty)$ at the linear rate*

$$(2.12) \quad \|d^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \left(\frac{\alpha(1-\beta)}{\beta(1-\alpha)}\right)^k \|e^0(\beta L, \cdot)\|_\infty,$$

$$(2.13) \quad \|e^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \left(\frac{\alpha(1-\beta)}{\beta(1-\alpha)}\right)^k \|d^0(\alpha L, \cdot)\|_\infty.$$

Proof. Since the errors d^k and e^k are both in the kernel of the heat operator they obtain, by the maximum principle, their maximum value on the boundary or on the initial line. On the initial line and the exterior boundary, both d^k and e^k vanish. Hence

$$\|d^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|e^{2k}(\beta L, \cdot)\|_\infty, \quad \|e^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|d^{2k}(\alpha L, \cdot)\|_\infty.$$

Using Lemma 2.3 the result follows. \square

2.2. Semidiscrete case. Consider the heat equation continuous in time, but discretized in space using a centered second-order finite difference scheme on a grid with n grid points and $\Delta x = \frac{L}{n+1}$. This gives the linear system of ODEs

$$(2.14) \quad \begin{aligned} \frac{\partial \mathbf{u}}{\partial t} &= A_{(n)} \mathbf{u} + \mathbf{f}(t), \quad t > 0, \\ \mathbf{u}(0) &= \mathbf{u}_0, \end{aligned}$$

where the $n \times n$ matrix $A_{(n)}$, the vector-valued function $\mathbf{f}(t)$, and the initial condition \mathbf{u}_0 are given by

$$(2.15) \quad A_{(n)} = \frac{1}{(\Delta x)^2} \begin{bmatrix} -2 & 1 & & 0 \\ 1 & -2 & \ddots & \\ & \ddots & \ddots & 1 \\ 0 & & 1 & -2 \end{bmatrix}, \quad \mathbf{f}(t) = \begin{pmatrix} f(\Delta x, t) + \frac{1}{(\Delta x)^2} g_1(t) \\ f(2\Delta x, t) \\ \vdots \\ f((n-1)\Delta x, t) \\ f(n\Delta x, t) + \frac{1}{(\Delta x)^2} g_2(t) \end{pmatrix}, \quad \mathbf{u}_0 = \begin{pmatrix} u_0(\Delta x) \\ \vdots \\ u_0(n\Delta x) \end{pmatrix}.$$

We note the following property of $A_{(n)}$ for later use: let $\mathbf{p} := (p_1, \dots, p_n)^T$, where $p_j := j$. Then

$$(2.16) \quad A_{(n)}\mathbf{p} = \left(0, \dots, 0, \frac{-(n+1)}{(\Delta x)^2} \right)^T.$$

Likewise, let $\mathbf{q} := (q_1, \dots, q_n)^T$ where $q_j := n + 1 - j$. Then

$$(2.17) \quad A_{(n)}\mathbf{q} = \left(\frac{-(n+1)}{(\Delta x)^2}, 0, \dots, 0 \right)^T.$$

We denote the i th component of a vector-valued function $\mathbf{v}(t)$ by $\mathbf{v}(i, t)$, and $\mathbf{v}(t) \geq \mathbf{u}(t)$ is understood componentwise. We now establish the discrete analogs of the maximum principle (Theorem 2.1) and Corollary 2.2.

THEOREM 2.5 (semidiscrete maximum principle). *Assume $\mathbf{u}(t)$ solves the semidiscrete heat equation (2.14) with $\mathbf{f}(t) = (f_1(t), 0, \dots, 0, f_2(t))^T$ and $\mathbf{u}(0) = (u_1(0), \dots, u_n(0))^T$. If $f_1(t)$ and $f_2(t)$ are nonnegative for $t \geq 0$ and $\mathbf{u}(i, 0) \geq 0$ for $i = 1, \dots, n$ then*

$$\mathbf{u}(t) \geq 0 \quad \forall t \geq 0.$$

Proof. We follow Varga’s proof in [20]. By Duhamel’s principle the solution $\mathbf{u}(t)$ is given by

$$(2.18) \quad \mathbf{u}(t) = e^{A_{(n)}t}\mathbf{u}(0) + \int_0^t e^{A_{(n)}(t-s)}\mathbf{f}(s)ds.$$

The key is to note that the matrix $e^{A_{(n)}t}$ contains only nonnegative entries. To see why write $A_{(n)} = -2I_{(n)} + J_{(n)}$, where $J_{(n)}$ contains only nonnegative entries and $I_{(n)}$ is the identity matrix of size $n \times n$. We get

$$e^{A_{(n)}t} = e^{-2I_{(n)}t}e^{J_{(n)}t} = e^{-2t}e^{J_{(n)}t} = e^{-2t} \sum_{l=0}^{\infty} \frac{J_{(n)}^l t^l}{l!},$$

where the last expression clearly has only nonnegative entries. Since the matrix exponential in (2.18) is applied only to vectors with nonnegative entries, it follows that $\mathbf{u}(t)$ cannot become negative. \square

COROLLARY 2.6. *The solution $\mathbf{u}(t)$ of the semidiscrete heat equation (2.14) with $\mathbf{f}(t) = (\frac{1}{(\Delta x)^2}g_1(t), 0, \dots, 0, \frac{1}{(\Delta x)^2}g_2(t))^T$ and $\mathbf{u}_0 \equiv 0$ satisfies the inequality*

$$(2.19) \quad \|\mathbf{u}(j, \cdot)\|_{\infty} \leq \frac{n+1-j}{n+1}\|g_1(\cdot)\|_{\infty} + \frac{j}{n+1}\|g_2(\cdot)\|_{\infty}, \quad 1 \leq j \leq n.$$

Proof. Consider $\tilde{\mathbf{u}}(t)$ solving

$$(2.20) \quad \begin{aligned} \frac{\partial \tilde{\mathbf{u}}}{\partial t} &= A_{(n)}\tilde{\mathbf{u}} + \tilde{\mathbf{f}}, & t > 0, \\ \tilde{\mathbf{u}}(j, 0) &= \frac{n+1-j}{n+1}\|g_1(\cdot)\|_{\infty} + \frac{j}{n+1}\|g_2(\cdot)\|_{\infty}, & 1 \leq j \leq n, \end{aligned}$$

with $\tilde{\mathbf{f}} = (\frac{1}{(\Delta x)^2}\|g_1(t)\|_{\infty}, 0, \dots, 0, \frac{1}{(\Delta x)^2}\|g_2(t)\|_{\infty})^T$. Using the properties (2.16) and (2.17) of $A_{(n)}$ and the linearity of (2.20) we find that the solution $\tilde{\mathbf{u}}$ of (2.20) does not depend on t and is given by the steady state solution

$$\tilde{\mathbf{u}}(j) = \frac{n+1-j}{n+1}\|g_1(\cdot)\|_{\infty} + \frac{j}{n+1}\|g_2(\cdot)\|_{\infty}, \quad 1 \leq j \leq n.$$




FIG. 2.2. Decomposition in the semidiscrete case.

The difference $\phi(j, t) := \tilde{\mathbf{u}}(j) - \mathbf{u}(j, t)$ satisfies the equation

$$\begin{aligned} \frac{\partial \phi}{\partial t} &= A_{(n)} \phi + \begin{pmatrix} \frac{1}{(\Delta x)^2} (\|g_1(\cdot)\|_\infty - g_1(t)) \\ 0 \\ \vdots \\ 0 \\ \frac{1}{(\Delta x)^2} (\|g_2(\cdot)\|_\infty - g_2(t)) \end{pmatrix}, \quad t > 0, \\ \phi(j, 0) &= \frac{n+1-j}{n+1} \|g_1(\cdot)\|_\infty + \frac{j}{n+1} \|g_2(\cdot)\|_\infty, \quad 1 \leq j \leq n, \end{aligned}$$

and hence by the discrete maximum principle $\phi(j, t) \geq 0$ for all $t > 0$ and $1 \leq j \leq n$.

Thus

$$\mathbf{u}(j, t) \leq \frac{n+1-j}{n+1} \|g_1(\cdot)\|_\infty + \frac{j}{n+1} \|g_2(\cdot)\|_\infty, \quad 1 \leq j \leq n.$$

Likewise, from $\psi(j, t) := \tilde{\mathbf{u}}(j) + \mathbf{u}(j, t)$, we get

$$\mathbf{u}(j, t) \geq - \left(\frac{n+1-j}{n+1} \|g_1(\cdot)\|_\infty + \frac{j}{n+1} \|g_2(\cdot)\|_\infty \right), \quad 1 \leq j \leq n.$$

Hence we can bound the modulus of \mathbf{u} by

$$|\mathbf{u}(j, t)| \leq \frac{n+1-j}{n+1} \|g_1(\cdot)\|_\infty + \frac{j}{n+1} \|g_2(\cdot)\|_\infty, \quad 1 \leq j \leq n.$$

Now the right-hand side does not depend on t , so we can take the supremum over t , which leads to the desired result. \square

We decompose the domain into two overlapping subdomains Ω_1 and Ω_2 as in Figure 2.2. We assume for simplicity that αL falls on the grid point $i = a$ and βL on the grid point $i = b$. We therefore have $a\Delta x = \alpha L$ and $b\Delta x = \beta L$. For notational convenience we define

$$\begin{aligned} \mathbf{f}^1(\mathbf{x}, y, z) &:= \left(\mathbf{x}(1) + \frac{y}{(\Delta x)^2}, \mathbf{x}(2), \dots, \mathbf{x}(b-2), \mathbf{x}(b-1) + \frac{z}{(\Delta x)^2} \right)^T, \\ \mathbf{f}^2(\mathbf{x}, y, z) &:= \left(\mathbf{x}(a+1) + \frac{y}{(\Delta x)^2}, \mathbf{x}(a+2), \dots, \mathbf{x}(n-1), \mathbf{x}(n) + \frac{z}{(\Delta x)^2} \right)^T. \end{aligned}$$

As in the continuous case, the solution $\mathbf{u}(t)$ of (2.14) can be obtained from the solutions $\mathbf{v}(t)$ on Ω_1 and $\mathbf{w}(t)$ on Ω_2 , which satisfy the equations

$$(2.21) \quad \begin{aligned} \frac{\partial \mathbf{v}}{\partial t} &= A_{(b-1)} \mathbf{v} + \mathbf{f}^1(\mathbf{f}(t), g_1(t), \mathbf{w}(b-a, t)), \quad t > 0, \\ \mathbf{v}(j, 0) &= \mathbf{u}_0(j), \quad 1 \leq j < b, \end{aligned}$$

and

$$(2.22) \quad \begin{aligned} \frac{\partial \mathbf{w}}{\partial t} &= A_{(n-a)} \mathbf{w} + \mathbf{f}^2(\mathbf{f}(t), \mathbf{v}(a, t), g_2(t)), & t > 0, \\ \mathbf{w}(j-a, 0) &= \mathbf{u}_0(j), & b \leq j \leq n. \end{aligned}$$

Applying the Schwarz iteration to (2.21) and (2.22) we obtain

$$\begin{aligned} \frac{\partial \mathbf{v}^{k+1}}{\partial t} &= A_{(b-1)} \mathbf{v}^{k+1} + \mathbf{f}^1(\mathbf{f}(t), g_1(t), \mathbf{w}^k(b-a, t)), & t > 0, \\ \mathbf{v}^{k+1}(j, 0) &= \mathbf{u}_0(j), & 1 \leq j < b, \end{aligned}$$

and

$$\begin{aligned} \frac{\partial \mathbf{w}^{k+1}}{\partial t} &= A_{(n-a)} \mathbf{w}^{k+1} + \mathbf{f}^2(\mathbf{f}(t), \mathbf{v}^k(a, t), g_2(t)), & t > 0, \\ \mathbf{w}^{k+1}(j-a, 0) &= \mathbf{u}_0(j), & b \leq j \leq n. \end{aligned}$$

Let $\mathbf{d}^k(t) := \mathbf{v}^k(t) - \mathbf{v}(t)$ and $\mathbf{e}^k(t) := \mathbf{w}^k(t) - \mathbf{w}(t)$ and consider the error equations

$$(2.23) \quad \begin{aligned} \frac{\partial \mathbf{d}^{k+1}}{\partial t} &= A_{(b-1)} \mathbf{d}^{k+1} + \mathbf{f}^1(0, 0, \mathbf{e}^k(b-a, t)), & t > 0, \\ \mathbf{d}^{k+1}(0) &= \mathbf{0}, \end{aligned}$$

and

$$(2.24) \quad \begin{aligned} \frac{\partial \mathbf{e}^{k+1}}{\partial t} &= A_{(n-a)} \mathbf{e}^{k+1} + \mathbf{f}^2(0, \mathbf{d}^k(a, t), 0), & t > 0, \\ \mathbf{e}^{k+1}(0) &= \mathbf{0}. \end{aligned}$$

The following lemma establishes convergence of the Schwarz iteration on the interface nodes of the subdomains in L^∞ . Using the discrete maximum principle, convergence in the interior then follows.

LEMMA 2.7. *On the interface gridpoints a and b the error of the Schwarz iteration decays at the rate*

$$(2.25) \quad \|\mathbf{d}^{k+2}(a, \cdot)\|_\infty \leq \frac{\alpha(1-\beta)}{\beta(1-\alpha)} \|\mathbf{d}^k(a, \cdot)\|_\infty,$$

$$(2.26) \quad \|\mathbf{e}^{k+2}(b, \cdot)\|_\infty \leq \frac{\alpha(1-\beta)}{\beta(1-\alpha)} \|\mathbf{e}^k(b, \cdot)\|_\infty.$$

Proof. By Corollary 2.6 we have

$$(2.27) \quad \|\mathbf{d}^{k+2}(j, \cdot)\|_\infty \leq \frac{j}{b} \|\mathbf{e}^{k+1}(b-a, \cdot)\|_\infty, \quad 1 \leq j < b,$$


and

$$(2.28) \quad \|\mathbf{e}^{k+1}(j, \cdot)\|_\infty \leq \frac{n+1-a-j}{n+1-a} \|\mathbf{d}^k(a, \cdot)\|_\infty, \quad 1 \leq j \leq b-a.$$

Evaluating (2.28) at $j = b-a$ and (2.27) at $j = a$ and combining the two we get

$$\|\mathbf{d}^{k+2}(a, \cdot)\|_\infty \leq \frac{a(n+1-b)}{b(n+1-a)} \|\mathbf{d}^k(a, \cdot)\|_\infty.$$

Now using $a\Delta x = \alpha L$, $b\Delta x = \beta L$, and $(n+1)\Delta x = L$ we get the desired result. The second inequality (2.26) is obtained similarly. \square

FIG. 3.1. Decomposition into N overlapping subdomains.

For any vector-valued function $\mathbf{h}(t)$ in $L^\infty(\mathbb{R}^+, \mathbb{R}^n)$ we define

$$\|\mathbf{h}(\cdot, \cdot)\|_{\infty, \infty} := \max_{1 < j < n} \|\mathbf{h}(j, \cdot)\|_{\infty}.$$

THEOREM 2.8. *The Schwarz iteration for the semidiscrete heat equation with two subdomains converges in $L^\infty(\mathbb{R}^+, \mathbb{R}^n)$ at the linear rate*

$$\begin{aligned} \|\mathbf{d}^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} &\leq \left(\frac{\alpha(1-\beta)}{\beta(1-\alpha)}\right)^k \|e^0(b-a, \cdot)\|_{\infty}, \\ \|e^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} &\leq \left(\frac{\alpha(1-\beta)}{\beta(1-\alpha)}\right)^k \|\mathbf{d}^0(a, \cdot)\|_{\infty}. \end{aligned}$$

Proof. By Corollary 2.6 we have

$$\|\mathbf{d}^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|e^{2k}(b-a, \cdot)\|_{\infty}, \quad \|e^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\mathbf{d}^{2k}(a, \cdot)\|_{\infty}.$$

Using Lemma 2.7 the result follows. \square

3. Arbitrary number of subdomains. We generalize the two-subdomain case described in section 2 to an arbitrary number of subdomains N . This leads to an algorithm which can be run in parallel. Subdomains with even indices depend only on subdomains with odd indices. Hence one can solve on all the even subdomains in parallel in one sweep, and then on all the odd ones in the next one. Boundary information is propagated in between sweeps.

Consider N subdomains Ω_i of Ω , $i = 1, \dots, N$, where $\Omega_i = [\alpha_i L, \beta_i L] \times [0, \infty)$ and $\alpha_1 = 0$, $\beta_N = 1$, and $\alpha_{i+1} < \beta_i$ for $i = 1, \dots, N-1$ so that all the subdomains overlap, as in Figure 3.1. We assume also that $\beta_i \leq \alpha_{i+2}$ for $i = 1, \dots, N-2$ so that domains which are not adjacent do not overlap. The solution $u(x, t)$ of (2.1) can be obtained as in the case of two subdomains by composing the solutions $v_i(x, t)$, $i = 1, \dots, N$, which satisfy the equations

$$(3.1) \quad \begin{aligned} \frac{\partial v_i}{\partial t} &= \frac{\partial^2 v_i}{\partial x^2} + f(x, t), & \alpha_i L < x < \beta_i L, \quad t > 0, \\ v_i(\alpha_i L, t) &= v_{i-1}(\alpha_i L, t), & t > 0, \\ v_i(\beta_i L, t) &= v_{i+1}(\beta_i L, t), & t > 0, \\ v(x, 0) &= u_0(x), & \alpha_i L \leq x \leq \beta_i L, \end{aligned}$$

where we have introduced for convenience of notation the two functions v_0 and v_{N+1} which are constant in x and satisfy the given boundary conditions, namely, $v_0(x, t) \equiv$




FIG. 3.2. Overlapping subdomains and corresponding error functions e_i .

$g_1(t)$ and $v_{N+1}(x, t) \equiv g_2(t)$. The system of equations (3.1), which is coupled through the boundary, can be solved using the Schwarz iteration. We get for $i = 1, \dots, N$

$$(3.2) \quad \begin{aligned} \frac{\partial v_i^{k+1}}{\partial t} &= \frac{\partial^2 v_i^{k+1}}{\partial x^2} + f(x, t), & \alpha_i L < x < \beta_i L, \quad t > 0, \\ v_i^{k+1}(\alpha_i L, t) &= v_{i-1}^k(\alpha_i L, t), & t > 0, \\ v_i^{k+1}(\beta_i L, t) &= v_{i+1}^k(\beta_i L, t), & t > 0, \\ v_i^{k+1}(x, 0) &= u_0(x), & \alpha_i L \leq x \leq \beta_i L, \end{aligned}$$

where again $v_0^k(t) \equiv g_1(t)$ and $v_{N+1}^k(t) \equiv g_2(t)$. Let $e_i^k := v_i^k(x, t) - v_i(x, t)$, $i = 1, \dots, N$, and consider the error equations (compare Figure 3.2)

$$(3.3) \quad \begin{aligned} \frac{\partial e_i^{k+1}}{\partial t} &= \frac{\partial^2 e_i^{k+1}}{\partial x^2}, & \alpha_i L < x < \beta_i L, \quad t > 0, \\ e_i^{k+1}(\alpha_i L, t) &= e_{i-1}^k(\alpha_i L, t), & t > 0, \\ e_i^{k+1}(\beta_i L, t) &= e_{i+1}^k(\beta_i L, t), & t > 0, \\ e_i^{k+1}(x, 0) &= 0, & \alpha_i L \leq x \leq \beta_i L, \end{aligned}$$

with $e_0^k(t) \equiv 0$ and $e_{N+1}^k(t) \equiv 0$.

For the following lemma, we need some additional definitions to facilitate the notation. We define $\alpha_0 = \beta_0 = 0$, $\alpha_{N+1} = \beta_{N+1} = 1$, and the constant functions $e_{-1} \equiv 0$ and $e_{N+2} \equiv 0$.

LEMMA 3.1. *The error e_i^{k+2} of the i th subdomain of the Schwarz iteration (3.3) decays on the interfaces $x = \beta_{i-1}L$ and $x = \alpha_{i+1}L$ at the rate*

$$(3.4) \quad \|e_i^{k+2}(\beta_{i-1}L, \cdot)\|_\infty \leq r_i r_{i+1} \|e_{i+2}^k(\beta_{i+1}L, \cdot)\|_\infty + r_i p_{i+1} \|e_i^k(\alpha_{i+1}L, \cdot)\|_\infty + p_i q_{i-1} \|e_i^k(\beta_{i-1}L, \cdot)\|_\infty + p_i s_{i-1} \|e_{i-2}^k(\alpha_{i-1}L, \cdot)\|_\infty$$

for $i = 2, \dots, N$ and

$$(3.5) \quad \|e_i^{k+2}(\alpha_{i+1}L, \cdot)\|_\infty \leq q_i r_{i+1} \|e_{i+2}^k(\beta_{i+1}L, \cdot)\|_\infty + q_i p_{i+1} \|e_i^k(\alpha_{i+1}L, \cdot)\|_\infty + s_i q_{i-1} \|e_i^k(\beta_{i-1}L, \cdot)\|_\infty + s_i s_{i-1} \|e_{i-2}^k(\alpha_{i-1}L, \cdot)\|_\infty$$

for $i = 1, \dots, N - 1$, where the ratios of the overlaps are given by

$$(3.6) \quad r_i = \frac{\beta_{i-1} - \alpha_i}{\beta_i - \alpha_i}, \quad p_i = \frac{\beta_i - \beta_{i-1}}{\beta_i - \alpha_i}, \quad q_i = \frac{\alpha_{i+1} - \alpha_i}{\beta_i - \alpha_i}, \quad s_i = \frac{\beta_i - \alpha_{i+1}}{\beta_i - \alpha_i}.$$

Proof. By Corollary 2.2 we have

$$(3.7) \quad \|e_i^{k+2}(x, \cdot)\|_\infty \leq \frac{x - \alpha_i L}{(\beta_i - \alpha_i)L} \|e_{i+1}^{k+1}(\beta_i L, \cdot)\|_\infty + \frac{\beta_i L - x}{(\beta_i - \alpha_i)L} \|e_{i-1}^{k+1}(\alpha_i L, \cdot)\|_\infty.$$

Note that the infinity norm of D and E equals 1. This can be seen, for example, for D by looking at the row sum of interior rows,

$$(3.11) \quad \begin{aligned} p_i s_{i-1} + p_i q_{i-1} + r_i p_{i+1} + r_i r_{i+1} &= p_i (s_{i-1} + q_{i-1}) + r_i (p_{i+1} + r_{i+1}) = p_i + r_i = 1, \\ s_i s_{i-1} + s_i q_{i-1} + q_i p_{i+1} + q_i r_{i+1} &= s_i (s_{i-1} + q_{i-1}) + q_i (p_{i+1} + r_{i+1}) = s_i + q_i = 1. \end{aligned}$$

The boundary rows, however, sum up to a value less than 1, namely

$$(3.12) \quad \begin{aligned} q_1 p_2 + q_1 r_2 &= q_1 (p_2 + r_2) = q_1 < 1, \\ p_{N-1} s_{N-2} + p_{N-1} q_{N-2} + r_{N-1} p_N &= p_{N-1} (s_{N-2} + q_{N-2}) + r_{N-1} p_N, \\ &= p_{N-1} + r_{N-1} p_N < 1, \\ s_{N-1} s_{N-2} + s_{N-1} q_{N-2} + q_{N-1} p_N &= s_{N-1} (s_{N-2} + q_{N-2}) + q_{N-1} p_N, \\ &= s_{N-1} + q_{N-1} p_N < 1. \end{aligned}$$

A similar result holds for the matrix E . Since the infinity norm of both D and E equals 1, convergence is not obvious at first glance. In the special case with two subdomains treated in section 2 the matrices E and D degenerate to the scalar $q_1 p_2$, which is strictly less than 1, and convergence follows. In the case of N subdomains the information from the boundary needs to propagate inward to the interior subdomains before the algorithm exhibits convergence. Hence we expect that the infinity norm of E and D raised to a certain power becomes strictly less than 1. We need the following lemmas to prove convergence.

LEMMA 3.2. *Let $\mathbf{r}(A) \in \mathbb{R}^p$ denote the vector containing the row sums of the $p \times p$ square matrix A . Then $\mathbf{r}(A^{n+1}) = A^n \mathbf{r}(A)$.*

Proof. Let $\mathbf{1} = (1, 1, \dots, 1)^T$. Then we have $\mathbf{r}(A) = A\mathbf{1}$ and hence $\mathbf{r}(A^{n+1}) = A^{n+1}\mathbf{1} = A^n A\mathbf{1} = A^n \mathbf{r}(A)$. \square

LEMMA 3.3. *Let A be a real $p \times q$ matrix with $a_{ij} \geq 0$ and B be a real $q \times r$ matrix with $b_{ij} \geq 0$. Define the sets $I_i(A) := \{k : a_{ik} > 0\}$ and $J_j(B) := \{k : b_{kj} > 0\}$. Then for the product $C := AB$ we have*

$$I_i(C) = \{k : I_i(A) \cap J_k(B) \neq \emptyset\}.$$

Proof. We have, since $a_{ik}, b_{kj} \geq 0$,

$$c_{ij} > 0 \iff \sum_{k=1}^q a_{ik} b_{kj} > 0 \iff \exists k \text{ s.t. } a_{ik} > 0 \text{ and } b_{kj} > 0 \iff I_i(A) \cap J_j(B) \neq \emptyset.$$

Hence for fixed i , $c_{ij} > 0$ if and only if $I_i(A) \cap J_j(B) \neq \emptyset$. \square

LEMMA 3.4. *D^k and E^k have strictly positive entries for all integer $k \geq \frac{N-1}{2}$.*

Proof. We show the proof for the matrix D ; the proof for E is similar. The row index sets $I_i(D)$ are given by

$$I_i(D) = \left\{ \begin{array}{ll} \left. \begin{array}{l} \{1, \dots, i+2\} \\ \{1, \dots, i+1\} \end{array} \right\} & \begin{array}{l} i \text{ even} \\ i \text{ odd} \end{array} \\ \left. \begin{array}{l} \{i-1, \dots, i+2\} \\ \{i-2, \dots, i+1\} \end{array} \right\} & \begin{array}{l} i \text{ even} \\ i \text{ odd} \end{array} \\ \left. \begin{array}{l} \{i-1, \dots, N-1\} \\ \{i-2, \dots, N-1\} \end{array} \right\} & \begin{array}{l} i \text{ even} \\ i \text{ odd} \end{array} \end{array} \right\} \begin{array}{l} 1 \leq i < 4, \\ 4 \leq i \leq N-3, \\ N-3 < i \leq N-1. \end{array}$$

The column index sets are given by

$$J_j(D) = \left\{ \begin{array}{ll} \left. \begin{array}{l} \{1, \dots, 3\} \\ \{j-1, \dots, j+2\} \\ \{j-2, \dots, j+1\} \end{array} \right\} & \begin{array}{l} j \text{ odd} \\ j \text{ even} \end{array} \\ \{N-2, N-1\} & \end{array} \right\} \begin{array}{l} 1 \leq j < 3, \\ 3 \leq j \leq N-2, \\ j = N-1. \end{array}$$

We are interested in the growth of the index sets $I_i(D^k)$ as a function of k . Once every index set contains all the numbers $1 \leq j \leq N-1$, the matrix D^k has strictly positive entries. We show that every multiplication with D enlarges the index sets $I_i(D^k)$ on both sides by two elements, as long as the elements 1 and $N-1$ are not yet reached. The proof is done by induction: for D^2 we have, using Lemma 3.3,

$$I_i(D^2) = \left\{ \begin{array}{ll} \{1, \dots, i+4\} & i \text{ even} \\ \{1, \dots, i+3\} & i \text{ odd} \end{array} \right\} \quad 1 \leq i < 6,$$

$$\left\{ \begin{array}{ll} \{i-3, \dots, i+4\} & i \text{ even} \\ \{i-4, \dots, i+3\} & i \text{ odd} \end{array} \right\} \quad 6 \leq i \leq N-5,$$

$$\left\{ \begin{array}{ll} \{i-3, \dots, N-1\} & i \text{ even} \\ \{i-4, \dots, N-1\} & i \text{ odd} \end{array} \right\} \quad N-5 < i \leq N-1.$$

Now suppose that for k we obtained the sets

$$I_i(D^k) = \left\{ \begin{array}{ll} \{1, \dots, i+2k\} & i \text{ even} \\ \{1, \dots, i+2k-1\} & i \text{ odd} \end{array} \right\} \quad 1 \leq i < 2+2k,$$

$$\left\{ \begin{array}{ll} \{i-2k+1, \dots, i+2k\} & i \text{ even} \\ \{i-2k, \dots, i+2k-1\} & i \text{ odd} \end{array} \right\} \quad 2+2k \leq i \leq N-2k-1,$$

$$\left\{ \begin{array}{ll} \{i-2k+1, \dots, N-1\} & i \text{ even} \\ \{i-2k, \dots, N-1\} & i \text{ odd} \end{array} \right\} \quad N-2k-1 < i \leq N-1.$$

Then for $k+1$ we have, applying Lemma 3.3 again,

$$I_i(D^{k+1}) = \left\{ \begin{array}{ll} \{1, \dots, i+2(k+1)\} & i \text{ even} \\ \{1, \dots, i+2(k+1)-1\} & i \text{ odd} \end{array} \right\} \quad 1 \leq i < 2+2(k+1),$$

$$\left\{ \begin{array}{ll} \{i-2(k+1)-1, \dots, i+2(k+1)\} & i \text{ even} \\ \{i-2(k+1), \dots, i+2(k+1)-1\} & i \text{ odd} \end{array} \right\} \quad 2+2(k+1) \leq i \leq N-2(k+1)-1,$$

$$\left\{ \begin{array}{ll} \{i-2(k+1)-1, \dots, N-1\} & i \text{ even} \\ \{i-2(k+1), \dots, N-1\} & i \text{ odd} \end{array} \right\} \quad N-2(k+1)-1 < i \leq N-1.$$

Hence every row index set $I_i(D^k)$ grows on both sides by 2 when D^k is multiplied by D , as long as the boundary numbers 1 and $N-1$ are not yet reached. Now the index set $I_1(D^k) = \{1, \dots, 2k\}$ has to grow most to reach the boundary number $N-1$, so we need for the number of iterations

$$k \geq \frac{N-1}{2}$$

for the matrix D^k to have strictly positive entries. \square

The infinity norm of a vector \mathbf{v} in \mathbb{R}^n and a matrix A in $\mathbb{R}^{n \times n}$ is defined by

$$\|\mathbf{v}\|_\infty := \max_{1 \leq j \leq n} |\mathbf{v}(j)|, \quad \|A\|_\infty := \max_{1 \leq i \leq n} \sum_{j=1}^n |A_{ij}|.$$

LEMMA 3.5. *For all $k > \frac{N}{2}$ there exists $\gamma = \gamma(k) < 1$ such that*

$$\|D^k\|_\infty \leq \gamma \quad \text{and} \quad \|E^k\|_\infty \leq \gamma.$$

Proof. We prove the result for D ; the proof for E is similar. We have from (3.11) and (3.12) that

$$r(D) = \begin{pmatrix} q_1 \\ 1 \\ \vdots \\ 1 \\ p_{N-1} + r_{N-1}p_N \\ s_{N-1} + q_{N-1}p_N \end{pmatrix}.$$

By Lemma 3.4 D^k has strictly positive entries for any $k \geq \frac{N}{2}$. Note also that $\|D^k\|_\infty \leq 1$ since $\|D\|_\infty \leq 1$. Now by Lemma 3.2 we have

$$\|D^{k+1}\|_\infty = \max_i r_i(D^{k+1}) = \max_i \sum_j D_{ij}^k r_j(D) < 1$$

since $D_{ij}^k > 0$ for all i, j , $\sum_j D_{ij}^k \leq 1$ for all i , $r_j(D) \in [0, 1]$ and $r_1(D) < 1$, $r_{N-1}(D) < 1$, and $r_N(D) < 1$. \square

Remark. It suffices for each row index set to reach one of the boundaries, either 1 or $N - 1$, for the infinity norm to start decaying. Hence it is enough that there are no more index sets $I_i(D^k)$ (compare the proof of Lemma 3.4) such that $2 + 2k \leq i \leq N - 1 - 2k$ so that the requirement $k \geq \frac{N-1}{2}$ can be relaxed to $k > \frac{N-3}{4}$.

We now fix some $k > \frac{N-3}{4}$ and set

$$(3.13) \quad \gamma := \max(\|D^k\|_\infty, \|E^k\|_\infty) < 1.$$

LEMMA 3.6. *The vectors ξ and η satisfy*

$$(3.14) \quad \|\xi^{2km}\|_\infty \leq \gamma^m \|\xi^0\|_\infty,$$

$$(3.15) \quad \|\eta^{2km}\|_\infty \leq \gamma^m \|\eta^0\|_\infty.$$

Proof. By induction on (3.8), using that the entries of D , E , ξ^k , and η^k are nonnegative, we get

$$\xi^{2km} \leq D^{km} \xi^0 \quad \text{and} \quad \eta^{2km} \leq E^{km} \eta^0.$$

Taking norms on both sides and applying Lemma 3.5 the result follows. \square

THEOREM 3.7. *The Schwarz iteration for the heat equation with N subdomains converges in the infinity norm in time and space. We have*

$$(3.16) \quad \max_{1 \leq 2i \leq N} \|e_{2i}^{2km+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \gamma^m \|\xi^0\|_\infty,$$

$$(3.17) \quad \max_{1 \leq 2i+1 \leq N} \|e_{2i+1}^{2km+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \gamma^m \|\eta^0\|_\infty.$$

Proof. We use again the maximum principle. Since the error e_i^k is in the kernel of the heat operator, by the maximum principle, e_i^k attains its maximum on the initial line or on the boundary. On the initial line e_i^k vanishes, therefore

$$\max_{1 \leq 2i \leq N} \|e_{2i}^{2km+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\xi^{2km}\|_\infty, \quad \max_{1 \leq 2i+1 \leq N} \|e_{2i+1}^{2km+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\eta^{2km}\|_\infty.$$

Using Lemma 3.6 the result follows. \square

Note that the bound for the rate of convergence in Theorem 3.7 is not explicit. This is unavoidable for the level of generality employed. But, if we assume for simplicity that the overlaps are all of the same size, then we can get more explicit rates




FIG. 4.1. Theoretical and measured decay rate of the error for two subdomains and three different sizes of the overlap.

THEOREM 3.10. *The Schwarz iteration for the heat equation with N subdomains that overlap at the same ratio $r \in (0, 0.5]$ converges in the infinity norm in time and space. Specifically, we have*

$$(3.18) \quad \max_{1 \leq 2i \leq N} \|e_{2i}^{2k}(\cdot, \cdot)\|_{\infty, \infty} \leq \left(1 - 4r(1-r) \sin^2 \frac{\pi}{2(N+1)}\right)^k \|\xi^0\|_2,$$

$$(3.19) \quad \max_{1 \leq 2i+1 \leq N} \|e_{2i+1}^{2k}(\cdot, \cdot)\|_{\infty, \infty} \leq \left(1 - 4r(1-r) \sin^2 \frac{\pi}{2(N+1)}\right)^k \|\eta^0\|_2.$$

Proof. From the proof of Theorem 3.7 we have

$$\max_{1 \leq 2i \leq N} \|e_{2i}^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\xi^{2k}\|_{\infty}, \quad \max_{1 \leq 2i+1 \leq N} \|e_{2i+1}^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\eta^{2k}\|_{\infty}.$$

Since the infinity norm is bounded by the spectral norm we get

$$\max_{1 \leq 2i \leq N} \|e_{2i}^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\xi^{2k}\|_2, \quad \max_{1 \leq 2i+1 \leq N} \|e_{2i+1}^{2k+1}(\cdot, \cdot)\|_{\infty, \infty} \leq \|\eta^{2k}\|_2.$$

Using Lemma 3.9 the result follows. \square

The results derived above for the continuous heat equation remain valid as in the two-subdomain case, when the heat equation is discretized. Details of this analysis can be found in [7].

4. Numerical experiments. We perform numerical experiments to measure the actual convergence rate of the algorithm for the example problem

$$(4.1) \quad \begin{aligned} \frac{\partial u}{\partial t} &= \frac{\partial^2 u}{\partial x^2} - e^{-(t-1)^2 - (x-\frac{1}{4})^2}, & 0 < x < 1, \quad 0 < t < 3, \\ u(0, t) &= e^{-2t}, & 0 < t < 3, \\ u(1, t) &= e^{-t}, & 0 < t < 3, \\ u(x, 0) &= 1, & 0 < x < 1. \end{aligned}$$

To solve the semidiscrete heat equation, we use the backward Euler method in time. The first experiment is done splitting the domain $\Omega = [0, 1] \times [0, 3]$ into the two




FIG. 4.2. Theoretical and measured decay rate of the error in the case of eight subdomains.

subdomains $\Omega_1 = [0, \alpha] \times [0, 3]$ and $\Omega_2 = [\beta, 1] \times [0, 3]$ for three pairs of values $(\alpha, \beta) \in \{(0.4, 0.6), (0.45, 0.55), (0.48, 0.52)\}$. Figure 4.1 shows the convergence of the algorithm on the grid point b for $\Delta x = 0.01$ and $\Delta t = 0.01$. The solid line is the predicted convergence rate according to Theorem 2.8, and the dashed line is the measured one. The measured error displayed is the difference between the numerical solution on the whole domain and the solution obtained from the domain decomposition algorithm. As initial guess for the iteration we used the initial condition constant in time. We also checked the robustness of the method by refining the time step and obtained similar results.

We solved the same problem (4.1) using eight subdomains which overlap by 35%. Figure 4.2 shows the decay of the infinity norm of ξ^k . The dashed line shows the measured decay rate, and the solid line the predicted one. Note that in the initial phase of the iteration the error stagnates, since information has to be propagated across domains.

Acknowledgments. We thank Gene Golub for showing us how to prove Lemma 3.8 and Olavi Nevanlinna, Morten Bjørhus, and Sigitas Keras for many interesting discussions.

REFERENCES

- [1] A. BELLEN AND M. ZENNARO, *The use of Runge-Kutta formulae in waveform relaxation methods*, Appl. Numer. Math., 11 (1993), pp. 95–114.
- [2] M. BJØRHUS, *On Domain Decomposition, Subdomain Iteration and Waveform Relaxation*, Dr. Ing. Thesis, Department of Mathematical Sciences, Norwegian Institute of Technology, University of Trondheim, Norway, 1995.
- [3] M. BJØRHUS, *A note on the convergence of discretized dynamic iteration*, BIT, 35 (1995), pp. 291–296.
- [4] K. BURRAGE, *Parallel and Sequential Methods for Ordinary Differential Equations*, Oxford University Press, New York, 1995.
- [5] J. R. CANNON, *The One-Dimensional Heat Equation*, Encyclopedia Math. Appl., Addison-Wesley, Reading, MA, 1984.

- [6] T. F. CHAN AND T. P. MATHEW, *Domain decomposition algorithms*, Acta Numerica, 3 (1994), pp. 61–143.
- [7] M. J. GANDER, *Space-Time Continuous Analysis of Waveform Relaxation for Partial Differential Equations*, Ph.D. Thesis, Scientific Computing and Computational Mathematics Program, Stanford University, Stanford, CA, 1997.
- [8] E. GILADI AND H. B. KELLER, *Space-time domain decomposition for parabolic problems*, SIAM J. Numer. Anal., submitted.
- [9] R. JELTSCH AND B. POHL, *Waveform relaxation with overlapping splittings*, SIAM J. Sci. Comput., 16 (1995), pp. 40–49.
- [10] E. LELARSMEE, A. E. RUEHLI, AND A. L. SANGIOVANNI-VINCENTELLI, *The waveform relaxation method for time-domain analysis of large scale integrated circuits*, IEEE Trans. CAD IC Syst., 1 (1982), pp. 131–145.
- [11] E. LINDELÖF, *Sur l'application des méthodes d'approximations successives à l'étude des intégrales réelles des équations différentielles ordinaires*, J. Math. Pures Appl., 10 (1894), pp. 117–128.
- [12] P. L. LIONS, *On the Schwarz Alternating Method*, Proceedings of the First International Symposium on Domain Decomposition Methods for Partial Differential Equations, Ecole Nationale des Ponts et Chaussées, Paris, France 1987.
- [13] CH. LUBICH AND A. OSTERMAN, *Multi-grid dynamic iteration for parabolic equations*, BIT, 27 (1987), pp. 216–234.
- [14] U. MIEKKALA AND O. NEVANLINNA, *Convergence of dynamic iteration methods for initial value problems*, SIAM J. Sci. Stat. Comput., 8 (1987), pp. 459–482.
- [15] O. NEVANLINNA, *Remarks on Picard-Lindelöf iterations*, Part I, BIT, 29 (1989), pp. 328–346, Part II, BIT, 29 (1989), pp. 535–562.
- [16] O. NEVANLINNA, *Domain decomposition and iterations in parabolic problems*, in Fourth International Symposium on Domain Decomposition Methods for Partial Differential Equations, Moscow, 1990, SIAM, Philadelphia, 1991.
- [17] D. O'LEARY AND R. E. WHITE, *Multi-Splittings of Matrices and Parallel Solution of Linear Systems*, SIAM J. Algebraic Discrete Methods, 6 (1985), pp. 630–640.
- [18] E. PICARD, *Sur l'application des méthodes d'approximations successives à l'étude de certaines équations différentielles ordinaires*, J. Math. Pures Appl., 9 (1893), pp. 217–271.
- [19] H. A. SCHWARZ, *Über einige Abbildungsaufgaben*, Ges. Math. Abh., 11 (1869), pp. 65–83.
- [20] R. S. VARGA, *Matrix Iterative Analysis*, Prentice-Hall, Englewood Cliffs, NJ, 1962.
- [21] E. ZAUDERER, *Partial Differential Equations of Applied Mathematics*, 2nd ed., John Wiley, New York, 1989.